

43



WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

PCT

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/165		(11) International Publication Number: WO 99/66919 (43) International Publication Date: 29 December 1999 (29.12.99)	
(21) International Application Number: PCT/US99/13064	(22) International Filing Date: 9 June 1999 (09.06.99)	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, FR, GE, GH, GM, GR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ZY, ARIPO patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, HE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GN, GW, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 09/100,284 19 June 1998 (19.06.98) US	(71) Applicant: McNEIL-PPC, INC. (US/US); Grandview Road, Skillman, NJ 08558 (US).	Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(72) Inventors: OIHANNESIAN, Lena, A.; 490 Mallard Circle, Blue Bell, PA 19422 (US); NADIG, David; 591 Weikel Road, Lansdale, PA 19446 (US); HIGGINS, John, D., III; 10 N. New Street, West Chester, PA 19380 (US); REY, Max; Riedenerstrasse 69, CH-8304 Wallisellen (CH); MARTELLUCCI, Stephen, A.; 216 Whitaker Avenue, Mont Clere, PA 19453 (US).	(74) Agents: CIAMPOCERO, Audley, A., Jr. et al.; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08903-7333 (US).		
(54) Title: PHARMACEUTICAL COMPOSITION CONTAINING A SALT OF ACETAMINOPHEN AND AT LEAST ONE OTHER ACTIVE INGREDIENT			
(57) Abstract Pharmaceutical composition comprising an alkali or alkaline-earth metal salt of acetaminophen and at least one other active ingredient selected from the group consisting of analgesics, decongestants, expectorants, antihistamines, gastrointestinal agents, diuretics, bronchodilators and mixtures thereof.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TM	Turkmenistan
BE	Belgium	GN	Guinea	ML	Mali	TR	Turkey
BF	Burkina Faso	GR	Greece	MR	Morocco	TT	Trinidad and Tobago
BG	Bulgaria	HU	Hungary	MT	Malta	UA	Ukraine
BJ	Benin	IE	Ireland	MX	Mexico	US	United States of America
BR	Brazil	IL	Israel	MY	Malaysia	UZ	Uzbekistan
BY	Belarus	IS	Iceland	NI	Nicaragua	VN	Viet Nam
CA	Canada	IT	Italy	NE	Niger	YU	Yugoslavia
CF	Central African Republic	JP	Japan	NL	Netherlands	ZW	Zimbabwe
CG	Congo	KG	Kyrgyzstan	NO	Norway		
CH	Switzerland	KR	Republic of Korea	NZ	New Zealand		
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	LR	Liberia	PT	Portugal		
CN	China	KZ	Kazakhstan	RU	Russian Federation		
CU	Cuba	LC	Shant Loka	SD	Sudan		
CZ	Czech Republic	LI	Liechtenstein	SE	Sweden		
DE	Germany	LU	Luxembourg	SG	Singapore		
DK	Denmark	LV	Latvia				
EE	Estonia	LR	Liberia				

- 1 -

- 2 -

PHARMACEUTICAL COMPOSITION CONTAINING A SALT OF ACETAMINOPHEN AND AT LEAST ONE OTHER ACTIVE INGREDIENT

This is a continuation-in-part of application Serial No. 08/987,210, filed

- 5 December 9, 1997, which is a continuation-in-part of application Serial No. 08/771,176, filed December 20, 1996, both of which are hereby incorporated by reference.

FIELD OF THE INVENTION

10 The present invention relates to salts of acetaminophen and, more particularly, to alkali metal and alkaline-earth metal salts of acetaminophen.

BACKGROUND OF THE INVENTION

- 15 Acetaminophen (APAP) is a well established therapeutic agent having both analgesic and antipyretic activity. Acetaminophen's relatively poor solubility in water and its bitter taste, however, make it difficult to formulate into to consumer acceptable oral dosage forms. Most commercially available acetaminophen oral dosage forms incorporate a taste masking coating on the acetaminophen particles or employ flavors and sweeteners to mask the bitter taste of the drug.

- 20 Other approaches for dealing with the solubility and taste of acetaminophen include the formation of amino acid esters of acetaminophen. I. M. Kovach in *Diss. Abst. Int. B* 1975, 36(2), 734-5 describes the synthesis of p-acetamidophenyl glycinate (APG), α -p-acetamidophenyl aspartate (AAPA) and β -p-acetamidophenyl aspartate (BAPA). These esters are reported to have a less bitter taste than acetaminophen. APG-HBr was five times more water soluble than acetaminophen, whereas BAPA-HCl was four times less water soluble than APAP.

It is also known that the formation of an appropriate salt of a hydrophobic compound, such as a lipophilic carboxylic acid, will usually improve the aqueous solubility of the compound. Sodium ibuprofen and sodium naproxen are examples of pharmaceutically active lipophilic carboxylic acids which have improved aqueous solubility in their salt form. These salts are typically formed by reacting the carboxylic acid with a strong base, such as sodium hydroxide or potassium hydroxide.

- 10 USSR Inventor's Certificate No. 629,209, published September 11, 1978, describes a method of preparing bis- $[\beta$ -(4-acetylaminophenoxy)ethyl] ether by reacting 4-acetylaminophenol with an alkaline agent, such as potassium hydroxide, in a solution of an organic solvent, such as dimethylformamide, followed by reacting the resulting solution of potassium phenolate with chlorex at the boiling point of the reaction mixture. The resulting ether is reported as being useful for the treatment of animals with helminthic diseases.

- 15 USSR Inventor's Certificate 1,803,833, published March 23, 1993, describes a method of preparing acetaminophen for fluorescence intensity measurements. The acetaminophen sample was prepared by first dissolving in isopropyl alcohol and then treating with an 8% solution of potassium hydroxide solution and chloroform at a KOH:chloroform volume ratio of 3-4. Heating was then carried out for 15-20 minutes at 70-80°C before the measurement of the sample's fluorescence intensity.

- 20 While both of the of the above-identified USSR Inventor's Certificates report the treatment of acetaminophen with potassium hydroxide, neither document reports the isolation of any potassium salt of acetaminophen.

- 25 M.S. Yu et al. in US Patent No. 5,360,615 discusses a pharmaceutical carrier system for enhancing the solubility of acidic, basic or amphoteric pharmaceuticals by partial ionization to produce a highly concentrated primarily non-aqueous

solution suitable for filling softgels or for two-piece encapsulation or tablet formation. The acetaminophen solution comprised 25-40% (wt.) of acetaminophen, 0.4-1.0 moles of hydroxide ion per mole of acetaminophen and 1-20% (wt.) water in polyethylene glycol. An exemplary concentrated solution of acetaminophen suitable for use as a softgel fill contained 1 equivalent APAP (35% by wt.), 1 equivalent potassium hydroxide, and the balance polyethylene glycol 600.

US Patent No. 5,273,759 to D.L. Simmons describes the addition of $Mg(OH)_2$ in solid form to tablets containing APAP.


Both Yu et al. and Simmons fail to report the isolation of any discrete salts of acetaminophen.

A need exists for isolated salts of acetaminophen with improved aqueous solubility and taste when compared to the conventional form of acetaminophen.

SUMMARY OF THE INVENTION

The present invention provides isolated salts of acetaminophen. The isolated salts are preferably the alkali metal and alkaline-earth metal salts of acetaminophen.

In a further aspect of the invention the isolated salts have the formula:



wherein n is 1 or 2, M is alkali metal when n is 1 and M is alkaline-earth metal when n is 2 and x is from 0 to about 10. These salts have been shown to have both improved aqueous solubility and a less bitter taste than the free acid form of acetaminophen. The invention also includes methods of making such salts.

The present invention also provides compositions comprising the isolated salts of acetaminophen and at least one other active ingredient selected from the group consisting of: analgesics, decongestants, expectorants, antitussives, antihistamines, diuretics, gastrointestinal agents, diuretics, bronchodilators, sleep-inducing agents, and mixtures thereof.

Another aspect of the invention relates to the method of administering such salts, alone or in combination with other active ingredients, to mammals in the need of an analgesic and/or antipyretic therapeutic agent. The present invention further relates to orally administerable dosage forms containing salts of acetaminophen, alone or in combination with such other active ingredients.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 is a plot the results of dissolution tests for tablets containing acetaminophen free acid and the isolated salts of acetaminophen.


Figure 2 is a plot of acetaminophen plasma concentrations versus time for the bioequivalency study in dogs described in Example VII.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Prior to the present invention there has been no reported isolation of any discreet salts (phenolates) of APAP. Furthermore, *in situ* solution characterization of any deprotonated APAP species has not been reported either. As used in the present invention, the "free acid" of acetaminophen means the protonated phenolic form of APAP.

The lack of discussion on APAP salts in the scientific literature may be due in part to the fact that the anionic form of APAP is stable in aqueous solution (pH > 11) for only a short period of time (< 24h). If the salt is not quickly isolated after formation, p-aminophenolate (PAP) can form and result in discoloration of the resulting product.

As used in the present invention, isolated salts of acetaminophen refers to salts of p-hydroxyacetanilide which are formed by the deprotonation of the phenolic proton of acetaminophen. The isolated salts are preferably the alkali metal and alkaline-earth metal salts of acetaminophen. In a further aspect of the invention the isolated salts have the formula:



wherein n is 1 or 2, M is alkali metal when n is 1 and M is alkaline-earth metal when n is 2 and x is from 0 to about 10.

The salts of APAP are prepared via a one step aqueous reaction of APAP with the desired mono or divalent metal hydroxide. Suitable mono or divalent metal hydroxides include sodium hydroxide, calcium hydroxide, lithium hydroxide, potassium hydroxide, magnesium hydroxide and cesium hydroxide. The molar ratio of hydroxide to acetaminophen is about 1:2 to about 10:1, preferably about 1:2 to about 1:1. The APAP and metal hydroxide are dissolved in water or a mixture of water and a water-miscible organic solvent, such as acetonitrile, methanol, isopropanol, ethanol or tetrahydrofuran. The crude reaction products are then recovered or isolated by precipitation upon the addition of a less polar water miscible solvent such as acetonitrile, ethanol or tetrahydrofuran. Alternatively, the crude product can be recovered or crystallized by cooling (0°C) or lyophilization of the reaction mixture. The recovery or isolation should generally be carried out as soon as the reaction product is formed so as to reduce the likelihood of product

discoloration due to the formation of PAP. The final product may be dried under vacuum.

The APAP salts of the present invention are also amenable to cation exchange reactions. For example, an aqueous slurry or solution of a monovalent metal salt of acetaminophen is contacted with a divalent metal cation whereby the anhydrous, divalent metal salt of acetaminophen is formed via a cation exchange reaction. The salt is then immediately recovered. Specifically, $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}_2\text{Ca}$ may be prepared by reacting an aqueous solution of $\text{C}_8\text{H}_9\text{NO}_2\text{Na}$ with 0.5 equivalent of calcium chloride (CaCl_2). After drying, the resulting $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}_2\text{Ca}$ was found to be anhydrous.

In addition to the anhydrous form, various hydration states of APAP salts can be prepared depending on the reaction conditions. These hydrated salts preferably have less than 10 moles of water per mole of APAP salt, and includes, for example, acetaminophen sodium pentahydrate, acetaminophen sodium hexahydrate, acetaminophen sodium heptahydrate, acetaminophen calcium dihydrate and acetaminophen lithium hexahydrate.

The aqueous solubility at 22°C of the APAP salts of the present invention is 490-540, 450-470 and 13 mg/mL for sodium, lithium and calcium, respectively. Accordingly, the sodium, lithium and calcium salts have solubilities equivalent to approximately 260-280, 250-270, and 10 mg/mL, respectively, of APAP free acid.

The APAP salts have significantly increased dissolution rates compared to the conventional free acid form of acetaminophen. In 0.1N hydrochloric acid using USP Dissolution Apparatus 2 (paddle speed: 50 rpm) at 37°C, the concentration of acetaminophen at 30 seconds was as follows:

30

- 7 -

APAP Form (Powder)	Mg/mL of APAP
Sodium Salt	0.30
Lithium Salt	0.32
Calcium Salt	0.20
Free Acid (control)	0.02

Figure 1 illustrates the tablet dissolution rates of the salts of the present invention. The sodium, lithium and calcium salts of APAP and the conventional form of APAP were each compressed into tablets and the dissolution rates were evaluated using the conditions described above. The dissolution media was assayed for acetaminophen in the free acid form. Figure 1 shows that the salts of the present invention have significantly higher acetaminophen dissolution rates than the conventional free acid.

The calcium and sodium salts of acetaminophen have been observed not to have the bitter properties of the conventional free acid form of acetaminophen. The calcium salt was almost tasteless, while the sodium salt was observed to be somewhat salty. The improved taste properties of the salts of the present invention will allow for acetaminophen oral dosage forms with improved taste to be formulated.

The onset of action of acetaminophen is believed to be hastened, relative to the free acid form, with the isolated salts of the present invention. The increase solubility of the salts of the present invention, results in faster peak acetaminophen plasma concentration. This property will potentially provide faster onset of action of the analgesic and/or antipyretic activity of acetaminophen.

The acetaminophen salts of the present invention may be administered to a mammal in a therapeutically effective amount, which is an amount that produces the desired therapeutic response upon oral administration, and can be readily determined by one skilled in the art. In determining such amounts, the particular compound being

- 8 -

administered, the bioavailability characteristics of the compound, the dose regime, the age and weight of the patient, and other factors must be considered. A typical unit dose orally administered to a human would range from about 80-1000 mg (APAP free acid basis).

The compositions and methods of the present invention may also preferably include at least one other active ingredient selected from the group consisting of analgesics, decongestants, expectorants, antitussives, antihistamines, gastrointestinal agents, diuretics, bronchodilators, sleep-inducing agents and mixtures thereof.

When the other active ingredient is selected from the group consisting of decongestants, expectorants, antitussives, antihistamines and mixtures thereof, the compositions are particularly useful for the treatment of cough, cold, cold-like and/or flu symptoms in mammals, such as humans. As used in the present invention, cold-like symptoms include coryza, nasal congestion, upper respiratory infections, allergic rhinitis, otitis, and sinusitis.

The analgesics useful in combination with the acetaminophen salts of this invention include acetyl salicylic acid, indomethacin, optically active isomers or racemates of ibuprofen, naproxen, flurbiprofen, carprofen, tiaprofenic acid, cicloprofen, ketoprofen, ketorolac, etodolac, indomethacin, sulindac, fenoprofen, diclofenac, piroxicam, benzydolmine, nabumetone, tramadol, codeine, oxycodone, hydrocodone, pharmaceutically acceptable salts thereof and mixtures thereof. Cyclooxygenase-2 (COX-2) inhibitors, such as flosulide, nimesulide, celecoxib, 5-(4-aminosulfonyl-3-fluorophenyl)-4-cyclohexyl-2-methyl-oxazole, meloxicam, nabumetone, etodolac, pharmaceutically acceptable salts thereof and mixtures thereof, may be used as an analgesic in the present invention.

The decongestants (sympathomimetics) suitable for use in the compositions of the present invention include pseudoephedrine, phenylpropanolamine,

phenylephrine, ephedrine, pharmaceutically acceptable salts thereof and mixtures thereof.

The expectorants (also known as mucolytic agents) preferred for use in the present invention include guaifenesin, glyceryl guaiacolate, terpin hydrate, ammonium chloride, N acetylcysteine, bromhexine, ambroxol, domiodol, 3-iodo-1,2-propanediol, pharmaceutically acceptable salts thereof and mixtures thereof.

The antitussives preferred for use in the present invention include those such as dextromethorphan, chlorphedianol, carbetapentane, caramiphen, noscapine, diphenhydramine, codeine, hydrocodone, hydromorphone, fominoben, benzonate, pharmaceutically acceptable salts thereof and mixtures thereof.

The antihistamines which may be employed include chlorpheniramine, brompheniramine, dexchlorpheniramine, dexbrompheniramine, triprolidine, doxylamine, tripelemamine, cyproheptadine, hydroxyzine, pyrilamine, azatadine, promethazine, acrivastine, astemizole, cetirizine, ketotifen, loratadine, temelastine, terfenadine, norastemizole, fexofenadine, pharmaceutically acceptable salts thereof and mixtures thereof.

Examples of gastrointestinal agents preferred for use in the present invention include anticholinergics, including: atropine, clidinium and dicyclomine; antacids, including: aluminum hydroxide, bismuth subsalicylate, bismuth subcitrate, calcium carbonate and magaldrate; anti-gas agents, including simethicone; H₂-receptor antagonists, including: cimetidine, famotidine, nizatidine and ranitidine; laxatives, including: phenolphthalein and casanthrol; gastroprotectants, including: sucralfate and sucralfate humid gel; gastrokinetic agents, including: metoclopramide and cisapride; proton pump inhibitors, including omeprazole and antidiarrheals, including: diphenoxylate and loperamide; pharmaceutically acceptable salts thereof and mixtures thereof.

The diuretics useful in the invention include caffeine and pamabrom. Also useful are bronchodilators such as terbutaline, aminophylline, pinephrine, isoprenaline, metaproterenol, bitoterol, theophylline, albuterol, pharmaceutically acceptable salts thereof and mixtures thereof.

Sleep-inducing agents suitable for use in the invention include melatonin, esazolam, zolpidem, promethazine, pharmaceutically acceptable salts thereof and mixtures thereof.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from nonorganic bases include sodium, potassium, lithium, ammonium, calcium, magnesium, ferrous, zinc, manganese, aluminum, ferric, manganese salts and the like. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, tertiary and quaternary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as triethylamine, tripropylamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, lysine, arginine, histidine, caffeine, procaine, N-ethylpiperidine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglycine, theobromine, purines, piperazine, piperidine, polyamine resins and the like.

As with the acetaminophen salts of the present invention, these other active ingredients are administered to a mammal in a therapeutically effective amount, which is an amount that produces the desired therapeutic response upon oral administration, and can be readily determined by one skilled in the art. In determining such amounts, the particular compound being administered, the bioavailability characteristics of the compound, the dose regime, the age and weight of the patient, and other factors must be considered. Many of these other active ingredients, as well as their acceptable dosage ranges are described in the following: U.S. Pat. No. 4,552,899 to Sunshine et

al., issued Nov. 12, 1985; U.S. Pat. No. 4,783,465 to Sunshine et al., issued Nov. 8, 1988; and U.S. Pat. No. 4,619,934 to Sunshine et al., issued Oct. 28, 1986, which are all incorporated by reference herein. Other antitussives, expectorants,

antihistamines, decongestants, and gastrointestinal agents suitable for use in the invention are described in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa., 18th ed. Chapters 39, 42, 43, 58 and 59 (1990), which is hereby incorporated by reference. These other active ingredients may be administered concomitantly as a combination product with the acetaminophen salt or they may be administered as separate products prior to or after the administration of the APAP

10 salt.

The acetaminophen salts of the present invention, alone or in combination with the other active ingredients, are generally administered orally in a solid dosage form.

Suitable solid preparations include as swallowable, chewable or fast dissolving tablets,

15 pills, capsules, caplets, powders, wafers, sachets, gelatin coated tablets, softgels and granules. In preparing solid dosage forms, the salt of acetaminophen, alone in combination with such other active ingredients, can be mixed with conventional solid

fillers or carriers, such as corn starch, talc, calcium phosphate, calcium sulphate, calcium stearate, magnesium stearate, stearic acid, sorbitol, microcrystalline cellulose,

20 mannitol, gelatin, natural or synthetic gums, such as carboxymethylcellulose, methylcellulose, alginate, dextran, acacia gum, karaya gum, locust bean gum and other conventional carriers. Additionally, other excipients such as diluents, binders, lubricants, disintegrants, colors and flavoring agents may be employed. The dosage

form can also be film coated. It may also be desirable to coat the acetaminophen salt and/or other active ingredients with a conventional, pharmaceutically acceptable

polymeric film prior to the preparation of the dosage form.

25 Conventional methods can be used for preparing the solid dosage forms of the present invention. Suitable techniques are described in *Remington's Pharmaceutical Sciences*, 18th Ed., Chapter 89 (1990) which is hereby incorporated by reference.

30

The following example illustrates a specific embodiment of the present invention. This invention, however, is not confined to the specific limitations set forth in this example but rather to the scope of the appended claims. Unless otherwise stated, the percentages and ratios given below are by weight.

5

EXAMPLE I

This Example discloses the preparation of acetaminophen sodium ($C_8H_9NO_2Na \cdot 6H_2O$).

10

30 mL 1N NaOH solution (0.030 mol) were added to a stirred suspension of 4.53 g (0.033 mol) acetaminophen in 25 mL water. After all solids dissolved, 200 mL acetonitrile was added while the solution was rapidly stirred. The resulting

15 white precipitate (9.15 g, 99% yield as the 6-hydrate) was collected on a frit, washed with tetrahydrofuran (THF) and dried at room temperature. 1H NMR (DMF d_4) δ 9.4 (s, 1H, NH), 7.1 (m, 2H, Ar-H), 6.3 (m, 2H, Ar-H), 1.96 (s, 3H, $CO-CH_3$); IR (cm $^{-1}$, KBr) 3421 (broad, OH), 1635 (sharp, CO), 1594 (sharp), 1534 (sharp), 1500 (sharp), 1279 (sharp); Combustion analysis calculated for $C_8H_9NO_2Na \cdot 6H_2O$: C 34.16, H

7.12, N 4.98, found C 34.05, H 6.96, N 5.00; Water content calculated for $C_8H_9NO_2Na \cdot 6H_2O$: 38%, Found: 38% (Karl Fischer); FAB mass spectral analysis

20 m/e calculated for $C_8H_9NO_2Na \cdot 6H_2O$: 173, found 174 (M + 1). The aqueous solubility at 22°C was 493 mg/mL.

EXAMPLE II

EXAMPLE II


25 This Example discloses the preparation of acetaminophen sodium ($C_8H_9NO_2Na \cdot 7H_2O$).

80g (2.00 mol) NaOH was dissolved in 400 mL water and added dropwise to a flask charged with 302g (2.00 mol) APAP dissolved in 2100 mL *i*-propanol, at 50°C with stirring. The solution was cooled to room temperature, whereupon an off-

white precipitate formed. The solids were filtered, washed with three 200 mL portions of *i*-propanol, and dried under a vacuum (500g, 84 % as the 7-hydrate). The ¹H NMR and IR spectra were identical to that of C₁₆H₁₆N₂O₄Na•6H₂O. Combustion analysis calculated for C₁₆H₁₆N₂O₄Na•7H₂O: C 32.11 H 7.41 N 4.68; Found: C 31.99, H 7.38, N 4.31; Water content calculated for C₁₆H₁₆N₂O₄Na•7H₂O: 42.1%; Found 42.7% (Karl Fischer). The aqueous solubility at 22°C was 541 mg/mL.

EXAMPLE III


This Example discloses the preparation of acetaminophen calcium



5g (0.033 mol) APAP and 1.22g (0.016 mol) Ca(OH)₂ were suspended in 200 mL water and the mixture was stirred for 4h, whereupon all solids went into solution. The solution was frozen in a bath of liquid nitrogen and lyophilized, leaving a light microcrystalline off-white solid (5.44g, 100% crude yield based on the hydrate X 2). ¹H NMR (DMF-d₄) δ 9.39 (s, 2H, NH), 7.15 (m, 4H, Ar), 6.80 (m, 4H, Ar), 2.10 (s, 6H, CO-CH₃). IR 3287 (broad), 1648 (sharp, C=O), 1594, 1541, 1500, 1279 (sharp) Combustion analysis calculated for C₁₆H₁₆N₂O₄Ca•2H₂O: C 51.05, H 5.36, N 7.45; Found: C 51.21, H 5.21, N 7.63. Water content calculated (Karl Fischer) for C₁₆H₁₆N₂O₄Ca•2H₂O: 9.6%; Found: 9.8%. The aqueous solubility at 22°C was 13 mg/mL.

EXAMPLE IV

This Example discloses the preparation of acetaminophen lithium



5g (0.033 mol) APAP was dissolved in 30 mL *i*-propanol/THF (1:3, degassed with argon). This solution was added rapidly to a flask charged with 1.38g of (0.033 mol) LiOH•H₂O dissolved in 20 mL water (argon degassed). The colorless solution was stored at 0° C for 16 h, whereupon white crystals formed. The crystals were filtered under argon, washed with THF and dried under a vacuum for 16 h (4.25g, 6 hydrate). ¹H NMR (DMF-d₃) δ 9.39 (s, 1H, NH), 7.15 (m, 2H, Ar-H), 6.80 (m, 2H, Ar-H), 2.10 (s, 3H, CO-CH₃); IR 3568 (sharp), 3402, 3243 (broad), 1672, 1618 (sharp), 1533, 1501, 1407, 1267, 1174 (sharp). Combustion analysis calculated for C₁₆H₁₆NO₃Li•6H₂O: C 36.23, H 7.60, N 5.28; Found: C 36.67, H 7.68, N 5.23; Water content calculated (Karl Fischer) for C₁₆H₁₆NO₃Li•6H₂O: 40.1%; Found: 38.4%. The aqueous solubility at 22°C was 455 mg/mL.

EXAMPLE V

This Example discloses an alternative preparation of acetaminophen lithium (C₁₆H₁₆NO₃Li•6H₂O).

Acetaminophen (15.1g; 0.1 mol), water, 90 mL and lithium hydroxide 1 N (100 mL, 0.1 mol) were placed in a 2 L beaker. After the solution became clear, acetonitrile (1500 mL) was added. The resulting white solids were filtered, washed with THF (ca. 500 mL) and dried at ambient leaving a dry white solid (23.0 g, 87% based on C₁₆H₁₆NO₃Li•6H₂O). ¹H NMR (DMF-d₃) δ 2.0 (s, 3H, CO-CH₃), 6.5 (m, 2H, Ar-H), 7.2 (m, 2H, Ar-H), 9.3 (s, 1H, Ac-NH-Ar); IR 3568 (sharp), 3402, 3243 (broad), 1672, 1618 (sharp), 1533, 1501, 1407, 1267, 1174 (sharp). Combustion analysis calculated for C₁₆H₁₆NO₃Li•6H₂O: C 36.23, H 7.60, N 5.28; Found: C 36.56, H 7.56, N 5.05. Water content calculated (Karl Fischer) for C₁₆H₁₆NO₃Li•6H₂O: 40.1%; Found: 40.0%. The aqueous solubility at 22°C was 472 mg/mL.

EXAMPLE VI

This Example discloses the preparation of an anhydrous acetaminophen calcium ($C_{16}H_{16}N_2O_4Ca$).

Acetaminophen (90.6g, 0.60 mol) was suspended in 135 mL water and a solution containing sodium hydroxide (24.0g, 0.6 mol) and 36mL water was added at 18-26°C over 30 min. To the resulting NaAPAP-slurry, a solution containing calcium chloride ($CaCl_2$) (44.1g, 0.3 mol) and 54 mL water was added at 20-25°C over 30 min. at room temperature. The reaction mixture was then heated to 60°C within 60 min. Immediately after reaching 60°C, the slurry was cooled to 20°C within 60 min. and stirred at 20°C for 30 min. The resulting $C_{16}H_{16}N_2O_4Ca$ (79g, 78%) was filtered off, washed with *i*-propyl alcohol (75 mL) and dried overnight at 80°C under vacuum. 1H NMR (D_2O) δ 7.01 (d, 8.4H), 6.57 (d, 8.4H), 2.06 (s, 6H, CO-CH₃). IR (cm⁻¹): 1651 (sharp, C=O), 1506, 1276, 854 (sharp). Combustion analysis calculated for $C_{16}H_{16}N_2O_4Ca$: C 55.65, H 4.7, N 8.23; Found: C 55.80, H 4.53, N 8.13.

EXAMPLE VII

A study was conducted in dogs to determine the bioavailability of acetaminophen sodium. The free acid form of acetaminophen was used as the control. Compressed cylindrical pellets having the following composition were prepared:

Acetaminophen Sodium - compressed neat (no excipients).

Control - 150 mg APAP, 30 mg microcrystalline cellulose, and 30 mg dextrates.

Eight male purebred beagles having a body weight at initial dosing of approximately 9 to 14 kg were used in the study. The dogs were fed PMI® Certified Canine Diet Meal No. 5007 and water, both *ad libitum*. The dogs were fasted overnight for approximately 12 hours prior to dosing and food was returned 4 hours after dosing.

The dogs were divided into two groups and each group was dosed with either acetaminophen sodium or the control (free acid APAP) pellets. A single dose equivalent to 300 mg of acetaminophen free acid was administered via an oral gavage using a stomach tube. Each dose was followed by 20 mL of water. After a period of one week, the each group was dosed again, but with the other form of acetaminophen. Twelve blood samples were collected from each dog on each dosing day (1 prior to dosing and 1 thereafter). The plasma was separated and tested for acetaminophen.

The following summarizes the pharmacokinetic measurements for acetaminophen:

Parameter	APAP Sodium	Control
AUC (ug-hr/mL)	31.4 ± 5.7	27.4 ± 6.1
C _{max} (ug/mL)	23.6 ± 4.2	19.4 ± 6.9
T _{max} (hr)	0.27 ± 0.1	0.60 ± 0.3
AUC = areas under the plasma concentration-time curve to the last quantifiable concentration. C _{max} = peak plasma concentration. T _{max} = peak time.		

Figure 2 is a plot of the acetaminophen plasma concentration-time curve. This Figure demonstrates that the acetaminophen salt of the present invention is absorbed faster than the free acid acetaminophen control. The faster T_{max} for the acetaminophen salt suggests faster onset of action of the analgesic and antipyretic activities relative to the free acid control.

- 17 -

EXAMPLE VIII

This Example discloses the preparation and testing of tablets containing anhydrous calcium acetaminophen (CaAPAP) and one other active ingredient selected from the group of chlorpheniramine maleate (CPM), dextromethorphan hydrobromide (DEX), diphenhydramine hydrochloride (DPH) and pseudoephedrine hydrochloride (PE). The target weight of the tablet (free APAP basis) was 325 mg. The following ingredients were used to make the tablets:

Ingredient	Formulation 1 (mg/Tab)	Formulation 2 (mg/Tab)	Formulation 3 (mg/Tab)	Formulation 4 (mg/Tab)
CaAPAP	368.23	368.23	368.23	368.23
CPM	2.00	-	-	-
DEX	-	15.00	-	-
DPH	-	-	25.00	-
PE	-	-	-	30.00
Microcrystalline Cellulose	520.77	507.77	497.77	492.77
(Avicel PH 200)				
SiO ₂ (Cab-O-Sil M5)	4.50	4.50	4.50	4.50
Mg Stearate NF	4.50	4.50	4.50	4.50

10

Appropriate amounts of these ingredients were weighed to make a 180 g batch. After sieving, the ingredients were combined and mixed using a PK Blender. The ingredients were then tableted using a single-punch Korsh tablet press. The weight, hardness, thickness and disintegration times were evaluated and are reported below. The dissolution of the CaAPAP was measured using USP Method II apparatus by monitoring the APAP concentration in gastric fluid(GF). The percent dissolution of APAP from the tablet formulations at 2 min. is also reported.

15


- 18 -

	Formulation 1	Formulation 2	Formulation 3	Formulation 4
Weight Range (mg)	917±6	900±4	907±9	913±4
Thickness range (mm)	5.72±0.02	5.65±0.03	5.72±0.02	5.56±0.02
Hardness range (kP)	7.9±0.1	9.1±0.3	7.1±1.1	8.8±0.5
Disintegration time (sec)	10 to 15	10 to 15	20	15 to 20
% dissolution of CaAPAP at 2 minutes in GF	100%	-	100%	100%

Various modifications can be made from the above-described embodiments without departing from the spirit and scope of the present invention.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising the isolated compound



wherein n is 1 or 2, M is alkali metal when n is 1 and M is alkaline-earth metal when n is 2 and x is from 0 to about 10, and at least one other active ingredient selected from the group consisting of analgesics, decongestants, expectorants, antitussives, antihistamines, gastrointestinal agents, diuretics, bronchodilators and mixtures thereof.

2. The composition of claim 2 wherein the alkali metal is selected from the group consisting of sodium, potassium, cesium and lithium.
3. The composition of claim 2 wherein the alkaline-earth metal is selected from the group consisting of calcium and magnesium.

4. The composition of claim 2 wherein the isolated compound is in a hydrated form.
5. The composition of claim 2 wherein the isolated compound is in an anhydrous form.

6. The composition of claim 2 wherein the analgesic is selected from the group consisting of acetyl salicylic acid, indomethacin, optically active isomers or racemates of ibuprofen, naproxen, flurbiprofen, carprofen, tiaprofenic acid, cicloprofen, ketoprofen, ketorolac, etodolac, indomethacin, sulindac, fenoprofen, diclofenac, piroxicam, benzydolmine, nabumetone, tramadol, codeine, oxycodone, hydrocodone, flosulide, nimesulide, celecoxib, 5-(4-aminosulfonyl-3-fluorophenyl)-

4-cyclohexyl-2-methylloxazole, meloxicam, nabumetone, etodolac, pharmaceutically acceptable salts thereof and mixtures thereof.

7. The composition of claim 2 wherein the decongestant is selected from the group consisting of pseudoephedrine, phenylpropanolamine, phenylephrine and ephedrine, pharmaceutically acceptable salts thereof and mixtures thereof.
8. The composition of claim 2 wherein the expectorant is selected from the group consisting of guaifenesin, glyceryl guaiacolate, terpin hydrate, ammonium chloride, N acetylcysteine and bromhexine, ambroxol, domitol, 3-iodo-1,2-propanediol, pharmaceutically acceptable salts thereof and mixtures thereof.
9. The composition of claim 2 wherein the antitussive is selected from the group consisting of dextromethorphan, chlorphedianol, carbetapentane, caramiphen, noscapine, diphenhydramine, codeine, hydrocodone, hydromorphone, fominoben, benzornate, pharmaceutically acceptable salts thereof and mixtures thereof.
10. The composition of claim 2 wherein the antihistamine is selected from the group consisting of chlorpheniramine, brompheniramine, dexchlorpheniramine, dexbrompheniramine, triprolidine, doxylamine, tripelemamine, cyproheptadine, hydroxyzine, pyrilamine, azelastine, promethazine, acrivastine, astemizole, cetirizine, keotifen, loratidine, temelastine, terfenadine, norastemizole, fexofenadine, pharmaceutically acceptable salts thereof and mixtures thereof.

- 21 -

11. The composition of claim 2 wherein the gastrointestinal agent is selected from the group consisting of atropine, clidinium, dicyclomine, aluminum hydroxide, bismuth subsalicylate, bismuth subcitrate, simethicone, calcium carbonate, magaldrate, cimetidine, famotidine, nizatidine, ranitidine, phenolphthalein, casanthrol, sucralate, sucralate humid gel, metoclopramide, cisapride, onaprazole, diphenoxylate, loperamide, pharmaceutically acceptable salts thereof and mixtures thereof.

12. The composition of claim 2 wherein the diuretic is selected from the group consisting of caffeine and pamabrom.

13. The composition of claim 2 wherein the bronchodilator is selected from the group consisting of terbutaline, aminophylline, pinephrine, isoprenaline, metaproterenol, bitoterol, theophylline, albuterol, pharmaceutically acceptable salts thereof and mixtures thereof.

14. The composition of claim 2 wherein the sleep-inducing agent is selected from the group consisting of melatonin, estazolam, zolpidem, promethazine, pharmaceutically acceptable salts thereof and mixtures thereof.

15. A method of eliciting an onset hastened analgesic or antipyretic response in a mammal, comprising the oral administration of the composition of claim 1.

16. The method of claim 15 wherein the alkali metal is selected from the group consisting of lithium, sodium, cesium and potassium.

17. The method of claim 15 wherein the alkaline-earth metal is selected from the group consisting of calcium and magnesium.

- 22 -

18. The method of claim 15 wherein the salt is in a hydrated form.

19. The method of claim 15 wherein the salt is in an anhydrous form.

FIG. 2





FIG. 1



INTERNATIONAL SEARCH REPORT

International Application No
PC./US 99/13064

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/165

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 23595 A (PROCTER & GAMBLE) 8 September 1995 (1995-09-08) page 6, line 23 -page 7, line 11; claims 1,7	1-14
A	WO 85 04589 A (SUNSHINE ABRAHAM; LASKA EUGENE M; STEGEL CAROLE E) 24 October 1985 (1985-10-24) page 2 -page 3	1-14
A	EP 0 396 404 A (SCHERING CORP) 7 November 1990 (1990-11-07) claim 1; examples 1-3	1,7,10
A	US 5 409 709 A (OZAMA KIYOTAKA ET AL) 25 Apr 11 1995 (1995-04-25) column 3 -column 4; claims 1,2	1,6,10,12

* Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" document published on or after the international filing date and not in conflict with the application but called to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is considered in combination with the prior art
- "Q" document referred to in an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "A" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

5 November 1999

12/11/1999

Authorized officer

Rufet, J

Name and mailing address of the ISA

European Patent Office, P.O. Box 5318, Paterden 2
D-69111 Heidelberg, Germany
Tel: (+31-70) 345-2840, Tx: 31 651 epo m,
Fax: (+31-70) 340-3016

INTERNATIONAL SEARCH REPORT

International Application No
PC./US 99/13064

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FR 2 751 875 A (SCR NEWPHARM) 6 February 1998 (1998-02-06) page 3, line 31 -page 4, line 41	1,6
A	PATENT ABSTRACTS OF JAPAN vol. 1997, no. 07, 31 July 1997 (1997-07-31) & JP 09 067256 A (TAISHO PHARMACEUT CO LTD), 11 March 1997 (1997-03-11) abstract	1,6
A	FR 2 278 324 A (BOTTU) 13 February 1976 (1976-02-13) page 1, line 5 -page 2, line 9	1
A	CHEMICAL ABSTRACTS, vol. 125, no. 11, 9 September 1996 (1996-09-09) Columbus, Ohio, US; abstract no. 142293w, page 1181; XP002059502 abstract & IN 172 949 A (RAMA, RAO INDIA)	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
WO 9523595		A		08-09-1995		US 5510389 A	
						AU 1935795 A	
						CA 2184365 A	
WO 8504589		A		24-10-1985		US 4552899 A	
						AT 72989 T	
						AU 2029195 A	
						AU 589554 B	
						AU 4120085 A	
						AU 7997998 A	
						CA 1258430 A	
						DE 3585495 A	
						EP 0180597 A	
						JP 2848556 B	
						JP 61501913 T	
						US 4749697 A	
						US 4839354 A	
						US 4749722 A	
						US 4749711 A	
						US 4749723 A	
						US 4749720 A	
						US 4749721 A	
						US 4783465 A	
						US 4920149 A	
						US 4840962 A	
						US 4871733 A	
						US 5025019 A	
						US 4619934 A	
						US 4738966 A	
EP 0396404		A		07-11-1990		US 4990535 A	
						AT 101517 T	
						AU 628986 B	
						AU 5664890 A	
						CA 2054752 A,C	
						DE 69006628 D	
						DE 69006628 T	
						DK 396404 T	
						EP 0471009 A	
						ES 2062355 T	
						HK 184896 A	
						JP 6006536 B	
						JP 4501425 T	
						KR 9411246 B	
						MX 9203278 A	
						WO 9013295 A	
						US 5100675 A	
US 5409709		A		25-04-1995		JP 5148139 A	
						JP 5246845 A	
						JP 5294829 A	
FR 2751875		A		06-02-1998		AU 3945197 A	
						CA 2233924 A	
						CZ 9801048 A	
						EP 0858329 A	
						WO 9805314 A	
						HU 9901893 A	

Form PCT/IS/A210 (patent family search) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
FR 2751875		A		13-02-1976		PL 326069 A	
JP 09067256		A		11-03-1997		NONE	
FR 2278324		A		13-02-1976		BE 831161 A	
						CH 595326 A	
						DE 2530535 A	
						GB 1514225 A	
						NL 7508210 A	
IN 172949		A		08-01-1994		NONE	

Form PCT/IS/A210 (patent family search) (July 1992)

INTERNATIONAL SEARCH REPORT		national application No. PCT/US 99/ 13064
Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)		
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:		
1. <input checked="" type="checkbox"/> Claims Nos.: 15-19 because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 15-19 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.		
2. <input type="checkbox"/> Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:		
3. <input type="checkbox"/> Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).		
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)		
This International Searching Authority found multiple inventions in this international application, as follows:		
1. <input type="checkbox"/> As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.		
2. <input type="checkbox"/> As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.		
3. <input type="checkbox"/> As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:		
4. <input type="checkbox"/> No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:		
Remark on Protest	<input type="checkbox"/> The additional search fees were accompanied by the applicant's protest.	<input type="checkbox"/> No protest accompanied the payment of additional search fees.